

# Chapter 2

## **Scope of the present thesis**

The current study was firstly performed to characterize how insomnia patients can be discriminated from people without sleep complaints with respect to behavior, brain function and brain structure. If differences exist, it is of importance to establish whether they represent risk factors that could have been present already before the onset of insomnia, or rather represent consequences of insomnia, i.e. induced by this chronic condition. We therefore planned our studies to allow for conclusions on inducibility and reversibility of possible differences. Thus, we evaluated the inducibility of effects by assessing subjects without sleep complaints both before and after enforcing the shallow sleep that is typical of insomnia, and we evaluated the reversibility of possible insomnia effects by assessing insomniacs both before and after treating them with non-pharmacological sleep therapy.

For two reasons, we focused on older adults, between 50 and 75 years of age. First, the prevalence of insomnia strongly increases with age<sup>4</sup>, making it the most relevant group to study. Second, we noticed that both experimental sleep deprivation and normal aging are associated most prominently with changes in prefrontal function. This observation, in combination with the age-related increase in sleep problems, led us to question to what extent sleep problems add to what is commonly referred to as 'age-related cognitive decline'. In other words: to what extent might this reflect 'poor-sleep-induced cognitive decline'?

To investigate whether the possible differences between insomnia patients and controls without sleep complaints normalize after well established non-pharmacological sleep therapy, CBT supported by chronobiological treatment was applied for 6 weeks, after which all assessments were performed once more. The group of controls without sleep complaints received selective slow wave suppression during two nights, either on their first or second session, which was randomly divided over participants.

On the basis of the current neuroimaging literature on sleep disruption, we expected to find prefrontal hypoactivation in insomnia, with compensatory activation in non-task related regions. We expected normalization of activation levels after sleep therapy, both on hypoactivation and on compensatory activation. On the basis of previous findings, we expected temporal and prefrontal regions to show less grey and white matter in insomnia patients compared to controls. We also expected the behavioral and brain activation outcomes of experimental selective slow wave suppression to be qualitatively comparable to insomnia findings, albeit possibly less strong.

In this thesis, we will try to answer the following questions:

- Given the equivocal neuropsychological test results of previous work, yet the pronounced subjective complaints, would it be possible to find insomnia-

- related performance differences on computerized tasks with high resolution reaction time sampling and subtle changes in task parameters?
- **Chapter 3** will focus on such a behavioral correlate of chronic insomnia, by measuring a simple and slightly more complex vigilance task before and after sleep therapy or waitlist control.
  - What are the brain activation patterns in older insomniacs performing two verbal fluency tasks that differ in complexity? If brain activation patterns differ from those of controls, can non-pharmacological sleep therapy normalize them? **Chapter 4** discusses how chronic insomnia patients differ from age-matched controls with respect to performance and brain activation on two verbal fluency tasks, and whether differences normalize after sleep therapy.
  - **Chapter 5** discusses structural brain correlates of chronic insomnia. Are there volumetric grey or white matter differences between chronic insomnia patients and age-matched controls? And if so, are they truly related to sleep complaints or rather confounds of factors like depression or age, that are known to affect regional brain volume?
  - Further 'state' factors of insomnia are discussed in **Chapter 6**, where the effects of double pulse transcranial magnetic stimulation are compared between groups and before and after sleep therapy. TMS was applied to test the hypothesis of changes in the balance between GABA-ergic inhibitory and glutamatergic excitatory neurons in cortical networks<sup>61</sup>. Do insomnia patients show normal levels of intracortical facilitation, normally induced by double pulse TMS?
  - Insomnia is known to be related to reduced slow wave sleep. To investigate the comparability between insomnia and experimentally induced shallow sleep, slow wave suppression was applied in older adults without sleep complaints, who performed the same tasks as insomnia patients. In **Chapter 7**, the performance on vigilance tasks is discussed before and after slow wave suppression.
  - **Chapter 8** discusses how selective slow wave suppression in normal sleeping controls not only affects memory performance but also affects hippocampal brain activation. The data are compared to similar findings after total sleep deprivation.

The general discussion in **Chapter 9** will integrate the findings to answer the questions whether insomnia is or is not characterized by alterations in performance and brain activation and whether these are reversible consequences of insomnia or rather irreversible possible risk factors for developing insomnia. This chapter includes a discussion on how our findings suggest directions for future research to reach a better understanding of the condition of chronic insomnia.

## REFERENCES

1. Lichstein KL, Durrence HH, Taylor DJ, Bush AJ, Riedel BW (2003): Quantitative criteria for insomnia. *Behavioral Research and Therapy* 41:427-445.
2. Buysse DJ (2008): Chronic insomnia. *American Journal of Psychiatry* 165:678-686.
3. Morphy H, Dunn KM, Lewis M, Boardman HF, Croft PR (2007): Epidemiology of insomnia: a longitudinal study in a UK population. *Sleep* 30:274-280.
4. Foley DJ, Monjan AA, Brown SL, Simonsick EM, Wallace RB, Blazer DG (1995): Sleep complaints among elderly persons: an epidemiologic study of three communities. *Sleep* 18:425-432.
5. Zhang B, Wing YK (2006): Sex differences in insomnia: a meta-analysis. *Sleep* 29:85-93.
6. Chesson A, Jr., Hartse K, Anderson WM, Davila D, Johnson S, Littner M, et al. (2000): Practice parameters for the evaluation of chronic insomnia. An American Academy of Sleep Medicine report. Standards of Practice Committee of the American Academy of Sleep Medicine. *Sleep* 23:237-241.
7. American Academy of Sleep M (2000): *International classification of sleep disorders, Revised: Diagnostic and coding manual*. Rochester: American Academy of Sleep Medicine.
8. Buysse DJ, Ancoli-Israel S, Edinger JD, Lichstein KL, Morin CM (2006): Recommendations for a standard research assessment of insomnia. *Sleep* 29:1155-1173.
9. Spielman AJ (1986): Assessment of Insomnia. *Clinical Psychology Review* 6:11-25.
10. Basta M, Chrousos GP, Vela-Bueno A, Vgontzas AN (2007): Chronic Insomnia and Stress System. *Sleep Medicine Clinics* 2:279-291.
11. Beaulieu-Bonneau S, LeBlanc M, Merette C, Dauvilliers Y, Morin CM (2007): Family history of insomnia in a population-based sample. *Sleep* 30:1739-1745.
12. Boomsma DI, van Someren EJ, Beem AL, de Geus EJ, Willemsen G (2008): Sleep during a regular week night: a twin-sibling study. *Twin Research and Human Genetics* 11:538-545.
13. Pallesen S, Nordhus IH, Kvale G, Havik OE, Nielsen GH, Johnsen BH, et al. (2002): Psychological characteristics of elderly insomniacs. *Scandinavian Journal of Psychology* 43:425-432.
14. Kales A, Caldwell AB, Soldatos CR, Bixler EO, Kales JD (1983): Biopsychobehavioral correlates of insomnia. II. Pattern specificity and consistency with the Minnesota Multiphasic Personality Inventory. *Psychosomatic Medicine* 45:341-356.
15. Kales A, Soldatos CR, Kales JD (1987): Sleep disorders: insomnia, sleepwalking, night terrors, nightmares, and enuresis. *Annals of Internal Medicine* 106:582-592.
16. Bonnet MH, Arand DL (2003): Insomnia, metabolic rate and sleep restoration. *Journal of Internal Medicine* 254:23-31.
17. Vgontzas AN, Bixler EO, Lin HM, Prolo P, Mastorakos G, Vela-Bueno A, et al. (2001): Chronic insomnia is associated with nyctohemeral activation of the hypothalamic-pituitary-adrenal axis: clinical implications. *Journal of Clinical Endocrinology & Metabolism* 86:3787-3794.
18. Rodenbeck A, Huether G, Ruther E, Hajak G (2002): Interactions between evening and nocturnal cortisol secretion and sleep parameters in patients with severe chronic primary insomnia. *Neuroscience Letters* 324:159-163.
19. Lombardo C, Baglioni C, Devoto A, Lucidi F, Violani C (2008): Sympathetic arousal and arousability predicts subsequent sleep quality. *International Journal of Psychophysiology* 69:224-224.
20. Stepanski E, Zorick F, Roehrs T, Young D, Roth T (1988): Daytime alertness in patients with chronic insomnia compared with asymptomatic control subjects. *Sleep* 11:54-60.

21. Edinger JD, Glenn DM, Bastian LA, Marsh GR, Daile D, Hope TV, et al. (2001): Sleep in the laboratory and sleep at home II: comparisons of middle-aged insomnia sufferers and normal sleepers. *Sleep* 24:761-770.
22. Bonnet MH, Arand DL (1995): 24-Hour metabolic rate in insomniacs and matched normal sleepers. *Sleep* 18:581-588.
23. Pavlova M, Berg O, Gleason R, Walker F, Roberts S, Regestein Q (2001): Self-reported hyperarousal traits among insomnia patients. *Journal of Psychosomatic Research* 51:435-441.
24. Landis CA, Savage MV, Lentz MJ, Brengelmann GL (1998): Sleep deprivation alters body temperature dynamics to mild cooling and heating not sweating threshold in women. *Sleep* 21:101-108.
25. Muentner NK, Watenpugh DE, Wasmund WL, Wasmund SL, Maxwell SA, Smith ML (2000): Effect of sleep restriction on orthostatic cardiovascular control in humans. *Journal of Applied Physiology* 88:966-972.
26. Kato M, Phillips BG, Sigurdsson G, Narkiewicz K, Pesek CA, Somers VK (2000): Effects of sleep deprivation on neural circulatory control. *Hypertension* 35:1173-1175.
27. Shaver JL, Johnston SK, Lentz MJ, Landis CA (2002): Stress exposure, psychological distress, and physiological stress activation in midlife women with insomnia. *Psychosomatic Medicine* 64:793-802.
28. Morin CM, Rodrigue S, Ivers H (2003): Role of stress, arousal, and coping skills in primary insomnia. *Psychosomatic Medicine* 65:259-267.
29. Bixler E, Vgontzas A, Lin H, Vela-Bueno A (2007): Menopause and hormone replacement therapy: Effects on objective sleep patterns. *Sleep* 30:A258-A259.
30. Lugaresi E, Zucconi M, Bixler EO (1987): Epidemiology of Sleep Disorders. *Psychiatric Annals* 17:446-&.
31. Fulda S, Schulz H (2001): Cognitive dysfunction in sleep disorders. *Sleep Medicine Reviews* 5:423-445.
32. Sateia MJ, Doghramji K, Hauri PJ, Morin CM (2000): Evaluation of chronic insomnia. *Sleep* 23:243-308.
33. Varkevisser M, Kerkhof GA (2005): Chronic insomnia and performance in a 24-h constant routine study. *Journal of Sleep Research* 14:49-59.
34. Szelenberger W, Niemcewicz S (2000): Severity of insomnia correlates with cognitive impairment. *Acta Neurobiologiae Experimentalis* 60:373.
35. Crenshaw MC, Edinger JD (1999): Slow-wave sleep and waking cognitive performance among older adults with and without insomnia complaints. *Physiology & Behavior* 66:485-492.
36. Orff HJ, Drummond SP, Nowakowski S, Perils ML (2007): Discrepancy between subjective symptomatology and objective neuropsychological performance in insomnia. *Sleep* 30:1205-1211.
37. Fang SC, Huang CJ, Yang TT, Tsai PS (2008): Heart rate variability and daytime functioning in insomniacs and normal sleepers: preliminary results. *Journal of Psychosomatic Research* 65:23-30.
38. Leger D, Massuel MA, Metlaine A (2006): Professional correlates of insomnia. *Sleep* 29:171-178.
39. Leger D, Guilleminault C, Bader G, Levy E, Paillard M (2002): Medical and socio-professional impact of insomnia. *Sleep* 25:625-629.
40. Simon GE, VonKorff M (1997): Prevalence, burden, and treatment of insomnia in primary care. *American Journal of Psychiatry* 154:1417-1423.
41. Godet-Cayre V, Pelletier-Fleury N, Le Vaillant M, Dinet J, Massuel MA, Leger D (2006): Insomnia and absenteeism at work. Who pays the cost? *Sleep* 29:179-184.

42. Daley M, Morin CM, LeBlanc M, Gregoire JP, Savard J (2009): The economic burden of insomnia: direct and indirect costs for individuals with insomnia syndrome, insomnia symptoms, and good sleepers. *Sleep* 32:55-64.
43. Ozminkowski RJ, Wang S, Walsh JK (2007): The direct and indirect costs of untreated insomnia in adults in the United States. *Sleep* 30:263-273.
44. Killgore WD, Grugle NL, Reichardt RM, Killgore DB, Balkin TJ (2009): Executive functions and the ability to sustain vigilance during sleep loss. *Aviation, Space, and Environmental Medicine* 80:81-87.
45. Nilsson JP, Soderstrom M, Karlsson AU, Lekander M, Akerstedt T, Lindroth NE, et al. (2005): Less effective executive functioning after one night's sleep deprivation. *Journal of Sleep Research* 14:1-6.
46. Harrison Y, Horne JA (2000): The impact of sleep deprivation on decision making: a review. *Journal of Experimental Psychology Applied* 6:236-249.
47. Killgore WD, Balkin TJ, Wesensten NJ (2006): Impaired decision making following 49 h of sleep deprivation. *Journal of Sleep Research* 15:7-13.
48. Chee MW, Choo WC (2004): Functional imaging of working memory after 24 hr of total sleep deprivation. *Journal of Neuroscience* 24:4560-4567.
49. Turner TH, Drummond SP, Salamat JS, Brown GG (2007): Effects of 42 hr of total sleep deprivation on component processes of verbal working memory. *Neuropsychology* 21:787-795.
50. Harrison Y, Horne JA (1998): Sleep loss impairs short and novel language tasks having a prefrontal focus. *Journal of Sleep Research* 7:95-100.
51. Vincent NK, Walker JR (2000): Perfectionism and chronic insomnia. *Journal of Psychosomatic Research* 49:349-354.
52. Lundh L, Broman, J, Hetta, J, Saboonchi, F (1994): Perfectionism and insomnia. *Scandinavian Journal of Behaviour Therapy* 23:3-18.
53. Nofzinger EA, Buysse DJ, Germain A, Price JC, Miewald JM, Kupfer DJ (2004): Functional neuroimaging evidence for hyperarousal in insomnia. *American Journal of Psychiatry* 161:2126-2128.
54. Staffen W, Mair A, Zauner H, Unterrainer J, Niederhofer H, Kutzelnigg A, et al. (2002): Cognitive function and fMRI in patients with multiple sclerosis: evidence for compensatory cortical activation during an attention task. *Brain* 125:1275-1282.
55. Feigin A, Ghilardi MF, Huang C, Ma Y, Carbon M, Guttman M, et al. (2006): Preclinical Huntington's disease: compensatory brain responses during learning. *Annals of Neurology* 59:53-59.
56. Smith MT, Perlis ML, Chengazi VU, Pennington J, Soeffing J, Ryan JM, et al. (2002): Neuroimaging of NREM sleep in primary insomnia: a Tc-99-HMPAO single photon emission computed tomography study. *Sleep* 25:325-335.
57. Smith MT, Perlis ML, Chengazi VU, Soeffing J, McCann U (2005): NREM sleep cerebral blood flow before and after behavior therapy for chronic primary insomnia: preliminary single photon emission computed tomography (SPECT) data. *Sleep Medicine* 6:93-94.
58. Lamarche CH, Ogilvie RD (1997): Electrophysiological changes during the sleep onset period of psychophysiological insomniacs, psychiatric insomniacs, and normal sleepers. *Sleep* 20:724-733.
59. Staner L, Cornette F, Maurice D, Viardot G, Le Bon O, Haba J, et al. (2003): Sleep microstructure around sleep onset differentiates major depressive insomnia from primary insomnia. *Journal of Sleep Research* 12:319-330.
60. Perlis ML, Kehr EL, Smith MT, Andrews PJ, Orff H, Giles DE (2001): Temporal and stagewise distribution of high frequency EEG activity in patients with primary and secondary insomnia and in good sleeper controls. *Journal of Sleep Research* 10:93-104.

61. Winkelman JW, Buxton OM, Jensen JE, Benson KL, O'Connor SP, Wang W, et al. (2008): Reduced brain GABA in primary insomnia: preliminary data from 4T proton magnetic resonance spectroscopy (1H-MRS). *Sleep* 31:1499-1506.
62. Drummond SP, Brown GG, Stricker JL, Buxton RB, Wong EC, Gillin JC (1999): Sleep deprivation-induced reduction in cortical functional response to serial subtraction. *Neuroreport* 10:3745-3748.
63. Yoo SS, Hu PT, Gujar N, Jolesz FA, Walker MP (2007): A deficit in the ability to form new human memories without sleep. *Nature Neuroscience* 10:385-392.
64. Drummond SP, Brown GG, Gillin JC, Stricker JL, Wong EC, Buxton RB (2000): Altered brain response to verbal learning following sleep deprivation. *Nature* 403:655-657.
65. Drummond SP, Meloy MJ, Yanagi MA, Orff HJ, Brown GG (2005): Compensatory recruitment after sleep deprivation and the relationship with performance. *Psychiatry Research* 140:211-223.
66. Drummond SP, Brown GG, Salamat JS, Gillin JC (2004): Increasing task difficulty facilitates the cerebral compensatory response to total sleep deprivation. *Sleep* 27:445-451.
67. Bell-McGinty S, Habeck C, Hilton HJ, Rakitin B, Scarmeas N, Zarahn E, et al. (2004): Identification and differential vulnerability of a neural network in sleep deprivation. *Cerebral Cortex* 14:496-502.
68. Chee MW, Chuah LY, Venkatraman V, Chan WY, Philip P, Dinges DF (2006): Functional imaging of working memory following normal sleep and after 24 and 35 h of sleep deprivation: Correlations of fronto-parietal activation with performance. *Neuroimage* 31:419-428.
69. Habeck C, Rakitin BC, Moeller J, Scarmeas N, Zarahn E, Brown T, et al. (2004): An event-related fMRI study of the neurobehavioral impact of sleep deprivation on performance of a delayed-match-to-sample task. *Cognitive Brain Research* 18:306-321.
70. Riemann D, Voderholzer U, Spiegelhalder K, Hornyak M, Buysse DJ, Nissen C, et al. (2007): Chronic insomnia and MRI-measured hippocampal volumes: a pilot study. *Sleep* 30:955-958.
71. Winkelman J, Benson K, Buxton O, Lyoo I, O'Connor S, Renshaw P (2009): Lack of Hippocampal Volume Differences in Primary Insomnia and Normal Sleeping Controls: An Mri Volumetric Study at 3 Tesla. *Sleep* 32:A260-A261.
72. Hall MH, Soreca I, Matthews KA, Kuller LH, Gianaros PJ (2009): Reported sleep duration and hippocampal grey matter volume in healthy women. *Sleep* 32:A3.
73. Jacobs GD, Pace-Schott EF, Stickgold R, Otto MW (2004): Cognitive behavior therapy and pharmacotherapy for insomnia: a randomized controlled trial and direct comparison. *Archives of Internal Medicine* 164:1888-1896.
74. Irwin MR, Cole JC, Nicassio PM (2006): Comparative meta-analysis of behavioral interventions for insomnia and their efficacy in middle-aged adults and in older adults 55+ years of age. *Health Psychology* 25:3-14.
75. Chesson AL, Jr., Anderson WM, Littner M, Davila D, Hartse K, Johnson S, et al. (1999): Practice parameters for the nonpharmacologic treatment of chronic insomnia. An American Academy of Sleep Medicine report. Standards of Practice Committee of the American Academy of Sleep Medicine. *Sleep* 22:1128-1133.
76. Riemann D, Perlis ML (2009): The treatments of chronic insomnia: A review of benzodiazepine receptor agonists and psychological and behavioral therapies. *Sleep Medicine Reviews*.
77. Morin CM, Colecchi C, Stone J, Sood R, Brink D (1999): Behavioral and pharmacological therapies for late-life insomnia: a randomized controlled trial. *Journal of the American Medical Association* 281:991-999.

78. Sivertsen B, Omvik S, Pallesen S, Bjorvatn B, Havik OE, Kvale G, et al. (2006): Cognitive behavioral therapy vs zopiclone for treatment of chronic primary insomnia in older adults: a randomized controlled trial. *Journal of the American Medical Association* 295:2851-2858.
79. Edinger JD, Wohlgemuth WK, Radtke RA, Marsh GR, Quillian RE (2001): Cognitive behavioral therapy for treatment of chronic primary insomnia: a randomized controlled trial. *Journal of the American Medical Association* 285:1856-1864.
80. Espie CA, Inglis SJ, Tessier S, Harvey L (2001): The clinical effectiveness of cognitive behaviour therapy for chronic insomnia: implementation and evaluation of a sleep clinic in general medical practice. *Behaviour Research and Therapy* 39:45-60.
81. Edinger JD, Wohlgemuth WK, Radtke RA, Coffman CJ, Carney CE (2007): Dose-response effects of cognitive-behavioral insomnia therapy: a randomized clinical trial. *Sleep* 30:203-212.
82. Spielman AJ, Saskin P, Thorpy MJ (1987): Treatment of chronic insomnia by restriction of time in bed. *Sleep* 10:45-56.
83. Morin CM, Hauri PJ, Espie CA, Spielman AJ, Buysse DJ, Bootzin RR (1999): Nonpharmacologic treatment of chronic insomnia. An American Academy of Sleep Medicine review. *Sleep* 22:1134-1156.
84. Edinger JD, Means MK (2005): Cognitive-behavioral therapy for primary insomnia. *Clinical Psychology Review* 25:539-558.
85. Riemersma-van der Lek RF, Swaab DF, Twisk J, Hol EM, Hoogendijk WJ, Van Someren EJ (2008): Effect of bright light and melatonin on cognitive and noncognitive function in elderly residents of group care facilities: a randomized controlled trial. *Journal of the American Medical Association* 299:2642-2655.
86. Van Someren EJW (2004): Sleep propensity is modulated by circadian and behavior-induced changes in cutaneous temperature. *Journal of Thermal Biology* 29:437-444.
87. Raymann RJ, Swaab DF, Van Someren EJ (2008): Skin deep: enhanced sleep depth by cutaneous temperature manipulation. *Brain* 131:500-513.